



BL

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A01N 25/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/60940</b> <b>(43) International Publication Date:</b> 19 October 2000 (19.10.00)
<b>(21) International Application Number:</b> PCT/US00/09568 <b>(22) International Filing Date:</b> 10 April 2000 (10.04.00) <b>(30) Priority Data:</b> 60/128,994 12 April 1999 (12.04.99) US <b>(71) Applicants:</b> DOW AGROSCIENCES LLC [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268-1054 (US). THE DOW CHEMICAL COMPANY [US/US]; 2030 Dow Center, Midland, MI 48674 (US). <b>(72) Inventors:</b> STROM, Robert, M.; 789 West Chippewa River Road, Midland, MI 48640 (US). PRICE, D., Claude; 2805 Canterbury Drive, Midland, MI 48667 (US). LUBETKIN, Steven, D.; 1135 Park Place, Zionsville, IN 46077 (US). <b>(74) Agent:</b> WILLIS, Reid, S.; Intellectual Property, P.O. Box 1967, Midland, MI 48641-1967 (US).		<b>(81) Designated States:</b> AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> AQUEOUS DISPERSIONS OF AGRICULTURAL CHEMICALS		
<b>(57) Abstract</b>  The bioavailability of a pesticide can be increased by formulating the pesticide as a stable aqueous dispersion with a particle mean diameter of not greater than 500 nm. Such a formulation has the further advantage of substantially reducing or eliminating the need for organic solvents. Moreover, the stable aqueous dispersion provides a means of preparing a one part formulation of a plurality of pesticides which would be otherwise unstable in each other's presence.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## AQUEOUS DISPERSIONS OF AGRICULTURAL CHEMICALS

The present invention relates to an aqueous dispersion of particles having pesticidal activity. In particular, the present invention relates to dispersions of micron- or sub-micron-sized particles.

5        The efficient use of pesticides is often restricted somewhat by their inherent poor water-solubility. Generally, these water-insoluble pesticides can be applied to a site in three ways, 1) as a dust, 2) as a solution in an organic solvent or a combination of water and one or more organic solvents, or 3) as an emulsion that is prepared by dissolving the product in an organic solvent, then dispersing the solution in water. All of these approaches have  
10 drawbacks. Application of a dust poses a health hazard and is inefficient. Solutions and emulsions that require an organic solvent are undesirable since the solvent usually serves no other purpose but to act as a carrier for the product. As such, the solvent adds an unnecessary cost to the formulation. It would be an advantage in the art, therefore, to provide a pesticide formulation that eliminates the need for organic solvents, on the one  
15 hand, but exhibits optimal availability to the site to which it is applied.

The present invention provides a stable, aqueous, pesticidal dispersion comprising from 1 to 60 percent by weight of the aqueous dispersion of a pesticide having a water solubility of less than 0.1 percent, having a crystalline phase, and having a melting point sufficiently high so as not to melt during milling, and from 1 to 30 percent by weight based  
20 on the weight of the pesticide of a surface active agent, wherein the dispersion has a volume mean diameter particle size of not greater than 500 nanometers (nm).

In a second aspect, the present invention is a method of preparing a stable aqueous pesticidal dispersion comprising the steps of:

- 25        a) mixing a pesticide with from 1 to 30 weight percent of a surface active agent and a sufficient amount of water to give an aqueous dispersion in the range of 1 to 60 weight percent, based on the weight of water and the pesticide; and
- b) reducing the volume mean diameter particle size of the pesticide to not greater than 500 nm using mechanical means and a grinding medium.

The present invention addresses a need in the art by providing aqueous dispersions of submicron-sized particles having high bioavailability and no or minimal amounts of an ancillary organic solvent.

5 The stable aqueous dispersion of the pesticide can be prepared by wet milling an aqueous dispersion of the pesticide in the presence of grinding media and a surface active agent. As used herein, the term "a surface active agent" means one or more surface active agents. The preparation is carried out in such a manner so as to produce a dispersion of non-agglomerating or interacting particles having a volume median diameter of not greater than 500 nm, more preferably not greater than 400 nm, most preferably not greater than  
10 200 nm, as measured by a Horiba model LA-700 particle size analyzer or an equivalent method. An example of such preparation can be found in U.S. Patent 5,145,684.

The pesticide is a solid at ambient temperature, and has a melting point that is sufficiently high that the pesticide does not melt through the milling process. The pesticide is preferably crystalline, and is substantially water insoluble. The term substantially water  
15 insoluble is used herein to refer to a pesticide that has a solubility in water of less than 0.1%, and most preferably less than 0.01%. It is understood that the water solubilities of many pesticides are pH-dependent, as a result of the functional groups they contain. Thus, pesticides with carboxylic acid groups or with sulfonamide or sulfonylurea groups, for example, may meet the solubility requirements at low pH but may be too soluble at high pH.  
20 The pH of the aqueous dispersion can be adjusted to ensure substantial insolubility of these pesticides.

Pesticides include herbicides, insecticides, and fungicides. Examples of classes of compounds that have herbicidal activity and meet the solubility, crystallinity and melting point requirements include, but are not restricted to, imidazolinones such as imazaquin, sulfonamide  
25 sulfonylureas such as chlorimuron-ethyl, triazolopyrimidine sulfonamides such as flumetsulam, aryloxyphenoxy propionates such as quizalofop ethyl, aryl ureas such as isoproturon and chlorotoluron, triazines such as atrazine and simazine, aryl carboxylic acids such as picloram, aryloxy alkanoic acids such as MCPA, chloroacetanilides such as metazachlor, dintroanilines such as oryzalin, pyrazoles such as pyrazolynate and diphenyl  
30 ethers such as bifenox.

Examples of classes of compounds that have insecticidal activity and meet the solubility, crystallinity and melting point requirements include, but are not restricted to, benzoyl ureas such as hexaflumuron, diacylhydrazines such as tebufenozide, carbamates such as carbofuran, pyrethroids such as *alpha*-cypermethrin, organophosphates such as phosmet, triazoles, and natural products such as spinosyns.

Examples of classes of compounds that have fungicidal activity and meet the solubility, crystallinity and melting point requirements include, but are not restricted to, morpholines such as dimethomorph, phenylamides such as benalaxyl, azoles such as hexaconazole, strobilurins such as azoxystrobin, phthalonitriles such as chlorothalonil and phenoxyquinolines such as quinoxyfen.

The surface active agent may be anionic, cationic, or nonionic, or combinations of cationic and nonionic or anionic and nonionic. Generally, and to a point, higher concentrations of surface active agent result in smaller particle size. Examples of suitable classes of surface active agents include, but are not limited to, anionics such as alkali metal fatty acid salts, including alkali metal oleates and stearates; alkali metal lauryl sulfates; alkali metal salts of diisooctyl sulfosuccinate; alkyl aryl sulfates or sulfonates, alkali metal alkylbenzene sulfonates, such as dodecylbenzene sulfonate; and alkali metal soaps; cationics such as long chain alkyl quaternary ammonium surfactants including cetyl trimethyl ammonium bromide; or nonionics such as ethoxylated derivatives of fatty alcohols, alkyl phenols, amines, fatty acids, fatty esters, mono-, di-, or triglycerides and various block copolymeric surfactants derived from ethylene oxide/propylene oxide. The concentration of the surface active agent is not less than 1%, and not more than 30% by weight, based on the total weight of the pesticide and the surface active agent. The surface active agent is preferably adsorbed onto the surface of the pesticide particle in accordance with U.S. Patent 5,145,684.

Examples of commercially available surface active agents include Atlox 4991 and 4913 surfactants (Uniqema), Morwet D425 surfactant (Witco), Pluronic P105 surfactant (BASF), Iconol TDA-6 surfactant (BASF), Kraftsperser 25M surfactant (Westvaco), Nipol 2782 surfactant (Stepan), Soprophor FL surfactant (Rhone-Poulenc), and Empicol LX 28 surfactant (Albright & Wilson).

The stable aqueous dispersions of the present invention can be prepared in the following manner. First, the pesticide is dispersed, preferably as a relatively coarse material having a particle size of less than 100  $\mu\text{m}$ . Such a degree of coarseness can be achieved, for example, by sieving. The coarse material is dispersed in an aqueous solution containing  
5 the surface active agent and preferably an antifoaming agent such as Antifoam B (Dow Corning) to form a premix. The premix can then be ground mechanically to reduce further the particle size of the pesticide. The concentration of the pesticide is in the range of from 1% to 60%, more preferably to 30%, by weight based on the weight of the pesticide and the water. Attrition time can vary widely depending on the mechanical means used and the  
10 choice and concentration of surfactant.

Examples of mechanical means to reduce the size of the agricultural product include a ball mill, a roller miller, an attritor mill, a sand mill, a bead mill, and a Cowles type mixer. The grinding media used for particle size reduction preferably include substantially  
15 spheroidal shaped particles having an average size of less than 3 mm. Examples of suitable media material include ZrO stabilized with magnesia, zirconium silicate, glass, stainless steel, polymeric beads, alumina, and titania, although nature of the material is not believed to be critical.

The particles are reduced in size at any suitable temperature where the agricultural product is stable. Typically, processing temperatures are not greater than the boiling point  
20 of water and below the melting point of the solid, but ambient temperature is preferred. The final volume mean diameter particle size is less than 500 microns, more preferably less than 400 nm, and most preferably less than 200 nm.

The stable aqueous dispersions are preferably diluted with water, then applied to a site in a pesticidally effective amount.

25 One of the advantages of the stable aqueous dispersion of the present invention is that it provides a means to prepare one-part formulations of different pesticides which are not only compatible with each other, but incompatible or unstable in each other's presence as well. For example, it may be desirable to combine a certain pesticide with a certain herbicide for a particular application but for the fact that the two (in solution, for example)  
30 react with each other faster than they can be applied to the desired site. However, in a stable aqueous dispersion, these different and incompatible pesticides can coexist, at least

temporarily, since they are shielded from each other from reacting rapidly, so that an end user can mix the incompatible pesticides together and apply them to a site before their efficacy is significantly diminished.

5 The fine particle size suspensions prepared according to the methods of the present invention, have enhanced biological efficacy as a result of their increased surface area as compared to suspensions of greater particle size. Enhanced biological efficacy is particularly important in cases where pesticide solubility is very low and thus, biological availability is restricted.

10 The following examples are for illustrative purposes only and are not intended to limit the scope of the invention. All percentages are by weight unless otherwise specified. All particle sizes (measured using a Horiba model LA-700 particle size analyzer) are in units of volume mean diameter.

Example 1 – Preparation of a Stable Aqueous Nano-Dispersion of a Triazole

15 1-Methyl- 3-(2-fluoro-6-chlorophenyl)-5-(3-methyl-4-bromothien-2-yl)-1H-1,2,4-triazole (1 g) was placed in a 20 mL polyethylene scintillation vial along with an aqueous phase (9 g) that was prepared by combining Pluronic P105 surfactant (3.33%), Morwet D425 surfactant (2.22%), deionized water (93.33%) and Antifoam B antifoaming agent (1.11%). About 10 g of 1/8" diameter stainless steel ball bearings were added, and the vial was then sealed, and placed a Retsch Model MM 2000 mill. The sample was milled for 90  
20 minutes and the particle size (pre-mill) was measured to be 1520 nm. A portion of the pre-mill (1 g) was removed and added to a fresh vial, at which time 1 drop of the antifoaming agent was added. Milling was continued for 3 hours, after which the particle size was determined to be 183 nm.

Example 2 – Preparation of a Stable Aqueous Nano-Dispersion Using Polymeric Beads

25 Example 1 was repeated except that styrene/divinylbenzene copolymer beads (475  $\mu$ m diameter) were used instead of stainless steel beads. After the pre-mill was subjected to 120 minutes of additional milling, the particle size was determined to be 148 nm.

Example 3 – Preparation of a Stable Aqueous Nano-Dispersion Using Alternative Surfactant Package

Example 1 was repeated except that the aqueous phase was prepared using Atlox 4991 surfactant (6%), Atlox 4913 surfactant (12%), deionized water (78%) and Antifoam B (4%). After the pre-mill was subjected to 270 minutes of additional milling, the particle size was determined to be 163 nm.

5 Example 4 - Preparation of a Stable Aqueous Nano-Dispersion of Spinosad

In this example, Spinosad insecticide was milled. Again, Example 1 conditions were used except that the dispersion was prepared using the surfactant package described in Example 3. After the pre-mill was subjected to 195 minutes of additional milling, the particle size was determined to be 297 nm.

10 Example 5 - Preparation of a Stable Aqueous Nano-Dispersion of Spinosad Using a Different Surfactant Package

The conditions of Example 1 were repeated except the Spinosad was used instead of the triazole. After the pre-mill was subjected to 315 minutes of additional milling, the particle size was found to be 281 nm.

15 Example 6 - Preparation of a Stable Aqueous Nano-Dispersion of Atrazine Herbicide

Atrazine herbicide (1 g) was placed in a 20 mL polyethylene scintillation vial along with an aqueous phase (9 g) prepared by combining Iconol TD-6 surfactant (0.5 g) deionized water (8 g) and Antifoam B (0.5 g). About 10 g of 1/8" stainless steel ball bearings were added, and the vial was sealed and placed on the mill. The particles were milled for 30 minutes, after which time a portion of the pre-mill (1 g) was removed and placed in a fresh vial along with fresh ball bearings (10 g) and 1 drop of Antifoam B. The vial was sealed and subjected to milling for an additional 105 minutes. Then Nipol 2782 surfactant (0.2 g) was added and milling was continued for an additional 2 hours. The particle size was measured to be 167 nm.

25 Example 7 - Preparation of a Stable Aqueous Nano-Dispersion of Epoxiconazole Herbicide

Epoxiconazole was placed in a 20 mL polyethylene scintillation vial along with an aqueous phase prepared from Soprophor FL surfactant (0.2 g), Empicol LX 28 (0.3 g), deionized water (8.45 g), and Antifoam B agent (0.05 g). About 10 g of 1/8" steel ball



bearings were added, and the vial was sealed and placed on the mill for 30 minutes, after which time, a portion of the pre-mill (1 g) was removed and placed in a fresh vial along with fresh ball bearings (10 g) and 1 drop of Antifoam B. The vial was sealed and subjected to milling for an additional 405 minutes. The particle size was measured to be 314 nm.

- 5 A correlation between particle and biological efficacy as demonstrated LC 50 is shown in the Table I below.

Table I

Two spotted spider mite	
VMD/ nm	LC <sub>50</sub>
0.404	15
0.372	11
0.332	7.6
0.163	4

- 10 As shown by the table, the amount required to kill 50% of the spider mites is substantially reduced at smaller VMD.

## WHAT IS CLAIMED IS:

1. A stable, aqueous pesticidal dispersion comprising from 1 to 60 percent by weight of the aqueous dispersion of a pesticide having a water solubility of less than 0.1 percent, having a crystalline phase, and having a melting point sufficiently high so as not to melt during milling, and from 1 to 30 percent by weight based on the weight of the pesticide of a surface active agent, wherein the dispersion has a volume mean diameter of not greater than 500 nm.
2. The pesticidal dispersion of Claim 1 wherein the pesticide is a herbicide selected from the group consisting of imidazolinones, sulfonylureas, triazolopyrimidine sulfonamides, aryloxyphenoxy propionates, aryl ureas, triazines, aryl carboxylic acids, aryloxy alkanoic acids, chloroacetanilides, dintroanilines, pyrazoles, and diphenyl ethers.
3. The pesticidal dispersion of Claim 1 wherein the pesticide is an insecticide selected from the groups consisting of benzoyl ureas, diacylhydrazines, carbamates, pyrethroids, organophosphates, triazoles, and natural products.
4. The pesticidal dispersion of Claim 1 wherein the pesticide is a fungicide selected from the group consisting of morpholines, phenylamides, azoles, strobilurins, phthalonitriles, and phenoxyquinolines.
5. The pesticidal dispersion of any of Claims 1 to 4 wherein the surface active agent is selected from the group consisting of alkali metal fatty acid salts, polyoxyethylene nonionics, alkali metal lauryl sulfates, quaternary ammonium surfactants, alkali metal alkylbenzene sulfonates, and alkali metal soaps.
6. The pesticidal dispersion of any of Claims 1 to 5 which has a volume mean diameter of not greater than 200 nm.
7. A method for controlling pests comprising the step of diluting the stable aqueous dispersion of any of Claims 1 to 6 with water and applying to a site a pesticidally effective amount of the diluted dispersion.

8. A method of preparing a stable aqueous pesticidal dispersion comprising the steps of:
- a) mixing a pesticide with from 1 to 30 weight percent of a surface active agent and a sufficient amount of water to give an aqueous dispersion in the range of 1 to 60 weight percent, based on the weight of water and the pesticide; and
  - b) reducing the volume mean diameter particle size of the pesticide to not greater than 500 nm using mechanical means and a grinding medium.
9. The method of Claim 8 wherein the mechanical means is a ball mill, a roller mill, an attritor mill, a sand mill, a bead mill, or a Cowles type mixer.
10. The method of either of Claims 8 or 9 wherein the grinding media are particles selected from the group consisting of ZrO stabilized with magnesia, zirconium silicate, glass, stainless steel, polymeric beads, alumina, and titania.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 00/09568

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N25/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	WO 99 48359 A (MULQUEEN PATRICK JOSEPH ;BANKS GRAHAM (GB); DIXON KEITH RODERICK ( ) 30 September 1999 (1999-09-30) page 1 -page 2 page 5 -page 10 claims 2-7,20 ---	1-10
Y	WO 95 07614 A (DOWELANCO ;MULQUEEN PATRICK JOSEPH (GB); BANKS GRAHAM (GB); LUBETK) 23 March 1995 (1995-03-23) page 1, paragraph 1 page 20, paragraph 3 examples 21-23 ---	1-10
Y	EP 0 499 299 A (STERLING WINTHROP INC) 19 August 1992 (1992-08-19) cited in the application page 2, line 50 -page 3, line 13; examples ---	1-10
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*B\* document member of the same patent family

Date of the actual completion of the international search

2 August 2000

Date of mailing of the international search report

11/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2260 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bertrand, F

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09568

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB 2 267 825 A (DOWELANCO) 22 December 1993 (1993-12-22) the whole document ----	1-10
X	US 4 832 950 A (TAKAYA SUSUMU ET AL) 23 May 1989 (1989-05-23) column 1 -column 3; examples ----	1-10
X	EP 0 683 979 A (AMERICAN CYANAMID CO) 29 November 1995 (1995-11-29) the whole document ----	1-10
Y	US 4 430 336 A (RENTZEA COSTIN ET AL) 7 February 1984 (1984-02-07) column 17, line 62 -column 18, line 4 -----	1-10

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09568

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9948359	A	30-09-1999	AU	2946299 A	18-10-1999
WO 9507614	A	23-03-1995	AU	691835 B	28-05-1998
			AU	7835594 A	03-04-1995
			BR	9407501 A	25-06-1996
			CA	2171848 A	23-03-1995
			EP	0719086 A	03-07-1996
			HU	74022 A,B	28-10-1996
			IL	110993 A	15-07-1998
			US	6074986 A	13-06-2000
			ZA	9407147 A	15-03-1996
EP 0499299	A	19-08-1992	US	5145684 A	08-09-1992
			AT	184202 T	15-09-1999
			AU	642066 B	07-10-1993
			AU	1014592 A	30-07-1992
			AU	654836 B	24-11-1994
			AU	1014792 A	30-07-1992
			CA	2059431 A	26-07-1992
			CA	2059432 A	26-07-1992
			DE	69229925 D	14-10-1999
			DE	69229925 T	17-02-2000
			EP	0498482 A	12-08-1992
			ES	2139586 T	16-02-2000
			FI	920321 A	26-07-1992
			FI	920322 A	26-07-1992
			HU	62462 A	28-05-1993
			HU	60635 A,B	28-10-1992
			IL	100754 A	16-10-1996
			IL	100755 A	08-12-1995
			JP	4317053 A	09-11-1992
			JP	4295420 A	20-10-1992
			KR	200061 B	15-06-1999
			MX	9200291 A	01-10-1992
			MX	9200292 A	01-10-1992
			NO	920333 A	27-07-1992
			NO	303668 B	17-08-1998
			NZ	241361 A	25-06-1993
			NZ	241362 A	25-06-1993
			SG	55104 A	21-12-1998
			RU	2074002 C	27-02-1997
			RU	2066553 C	20-09-1996
			US	5451393 A	19-09-1995
			US	5494683 A	27-02-1996
			US	5552160 A	03-09-1996
			US	5399363 A	21-03-1995
			US	5318767 A	07-06-1994
GB 2267825	A	22-12-1993	NONE		
US 4832950	A	23-05-1989	JP	1499790 C	29-05-1989
			JP	61057503 A	24-03-1986
			JP	63052002 B	17-10-1988
			JP	1011610 B	27-02-1989
			JP	1536817 C	21-12-1989
			JP	61151111 A	09-07-1986
			JP	1584937 C	31-10-1990
			JP	2011563 B	14-03-1990

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/09568

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4832950 A		JP 61152627 A	11-07-1986
		AT 69158 T	15-11-1991
		DE 3584595 A	12-12-1991
		EP 0173259 A	05-03-1986
		HK 41593 A	07-05-1993
		PH 23096 A	10-04-1989
		SG 14093 G	16-04-1993
EP 0683979 A	29-11-1995	US 5496845 A	05-03-1996
		AT 182048 T	15-07-1999
		AU 707944 B	22-07-1999
		AU 2019695 A	30-11-1995
		BG 62506 B	31-01-2000
		BG 99665 A	29-02-1996
		BR 9502522 A	09-04-1996
		CA 2149969 A	26-11-1995
		CN 1114134 A	03-01-1996
		CZ 9501259 A	13-12-1995
		DE 69510729 D	19-08-1999
		DE 69510729 T	04-11-1999
		EG 20392 A	28-02-1999
		ES 2133672 T	16-09-1999
		GR 3031104 T	31-12-1999
		HU 73158 A	28-06-1996
		JP 8040809 A	13-02-1996
		NO 952061 A	27-11-1995
		NZ 272184 A	24-03-1997
		PL 308737 A	27-11-1995
		RO 112684 A	30-12-1997
		SI 683979 T	31-10-1999
		SK 68795 A	06-12-1995
		TR 28583 A	04-11-1996
		ZA 9504215 A	22-01-1996
US 4430336 A	07-02-1984	DE 2948704 A	11-06-1981
		AT 2510 T	15-03-1983
		AU 6503180 A	11-06-1981
		BR 8007690 A	09-06-1981
		CA 1153002 A	30-08-1983
		CS 215143 B	30-07-1982
		DE 3062078 D	24-03-1983
		DK 515180 A	05-06-1981
		EP 0029996 A	10-06-1981
		HU 185622 B	28-03-1985
		IE 50726 B	25-06-1986
		IL 61515 A	29-02-1984
		JP 56092851 A	27-07-1981
		NZ 195740 A	07-12-1982
		PL 228213 A	07-08-1981
		SU 1060095 A	07-12-1983
		ZA 8007539 A	30-12-1981